

MATHEMATICS – III
(Common to EEE, ECE and EIE)

Time: 3 hours

Max. Marks: 70

PART – A
(Compulsory Question)

- 1 Answer the following: (10 X 02 = 20 Marks)
- Define Gamma function and evaluate $\int_0^{\infty} e^{-x^2} dx$.
 - Express $\int_0^1 \frac{dx}{\sqrt{1-x^4}}$ in terms of Beta function.
 - Express $\cos \theta$ and $\sin \theta$ in terms of Bessel function.
 - Prove that $P_n^1(1) = \frac{1}{2} n(n+1)$
 - Write the C – R equations in both Cartesian and Polar co-ordinates.
 - Find the fixed points of the transformation: $\omega = \frac{2i-6z}{iz-3}$.
 - State Cauchy's integral theorem.
 - Define pole of a complex function with example.
 - State Cauchy's residue theorem.
 - Evaluate $\int_C \frac{5z-2}{z(z-1)} dz$ where $C : |z| = 2$.

PART – B
(Answer all five units, 5 X 10 = 50 Marks)

UNIT – I

- 2 Show that $\beta(m, n) = 2 \int_0^{\pi/2} \sin^{2m-1} \theta \cos^{2n-1} \theta$ and deduce that $\int_0^{\pi/2} \sin^n \theta d\theta = \int_0^{\pi/2} \cos^n \theta d\theta = \frac{\Gamma(\frac{n+1}{2})\sqrt{\pi}}{2\Gamma(\frac{n+2}{2})}$.

OR

- 3 Find the power series solution of the equation $y'' + xy' + y = 0$ in powers of x .

UNIT – II

- 4 Prove that $P_n(x) = \frac{1}{2^n n!} \frac{d^n}{dx^n} (x^2 - 1)^n$ and hence express $2x^2 - 4x + 2$ as Legendre polynomial.

OR

- Prove that $\frac{d}{dx} [xJ_n(x)J_{n+1}(x)] = x[J_n^2(x) - J_{n+1}^2(x)]$.
- Prove that $J_{n+1}(x) = \frac{n}{x} J_n(x) - J_n'(x)$.

UNIT – III

- Determine P such that the function: $f(z) = \frac{1}{2} \log(x^2 + y^2) + i \tan^{-1} \left(\frac{yx}{x^2 - y^2} \right)$ be an analytic function.
- Find the analytic function, whose real part is $u = e^x [(x^2 - y^2) \cos y - 2xy \sin y]$.

OR

- Show that the function $\omega = \frac{4}{z}$ transform the straight line $x = c$ in the z -plane into a circle in the ω - plane.
- Find the bilinear transformation that maps the points $(0, i, 1)$ into the points $(-1, 0, 1)$.

UNIT – IV

- 8 Generate z^2 along the straight line OM and along the path OLM where 'O' is the origin, L is the point $z = 3$ and M is $z = 3 + i$ and hence establish the Cauchy's integral theorem.

OR

- Obtain the Taylor's series to represent the function $\frac{z^2-1}{(z+2)(z+3)}$ in the region $|z| < 2$.
- Expand $f(z) = \frac{e^{2z}}{(z-1)^3}$ about $z = 1$ as Laurent's series. Also find the region of convergence.

UNIT – V

- 10 Show that $\int_0^{2\pi} \frac{d\theta}{a+b \sin \theta} = \frac{2\pi}{\sqrt{a^2-b^2}}$, $a > b > 0$ using Residue theorem.

OR

- 11 Prove that $\int_{-\infty}^{\infty} \frac{x^2 dx}{(x^2+a^2)(x^2+b^2)} = \frac{\pi}{a+b}$ ($a > 0, b > 0, a \neq b$)
