

R16

Code No: 131AA

JAWAHARLAL NEHRU TECHNOLOGICAL UNIVERSITY HYDERABAD

B.Tech I Year I Semester Examinations, May - 2018

MATHEMATICS-I

(Common to CE, EEE, ME, ECE, CSE, EIE, IT, MCT, ETM, MMT, AE, MIE, PTM, CEE, MSNT)

Time: 3 hours

Max. Marks: 75

Note: This question paper contains two parts A and B.

Part A is compulsory which carries 25 marks. Answer all questions in Part A.

Part B consists of 5 Units. Answer any one full question from each unit. Each question carries 10 marks and may have a, b, c as sub questions.

PART-A

(25 Marks)

- 1.a) Find an integrating factor for the following equation  $\frac{dy}{dx} = e^{2x} + y - 1$ . [2]
- b) Find the solution of  $\frac{dy}{dx} = -\frac{x}{y}$  at  $x=1$  and  $y=\sqrt{3}$ . [3]
- c) Find the value of  $\alpha$  such that the vectors  $(1, 1, 0)$ ,  $(1, \alpha, 0)$  and  $(1, 1, 1)$  are linearly dependent. [2]
- d) Determine whether the system of equations is consistent 
$$\begin{aligned} 2x - 3y + 5z &= 1 \\ 3x + y - z &= 2 \\ x + 4y - 6z &= 1 \end{aligned}$$
 [3]
- e) If  $\lambda$  is the Eigen value of a matrix A then derive the Eigen value of (adjoint A). [2]
- f) Taking A as a  $2 \times 2$  matrix show that the Eigen values of A = the trace of A. [3]
- g) If  $u = x^y$  show that  $\frac{\partial^3 u}{\partial x^2 \partial y} = \frac{\partial^3 u}{\partial x \partial y \partial x}$ . [2]
- h) Find the stationary values of  $xy(a - x - y)$ . [3]
- i) Eliminate the arbitrary function  $f$  from the equation and form the partial differential equation  $z = xy + f(x^2 + y^2)$ . [2]
- j) Eliminate the constants  $a$  and  $b$  from the equation:  $z = (y + a)(x + b)$ . [3]

PART-B

(50 Marks)

- 2.a) Solve the Following differential equations:  
 $y'' - 2y' + y = te' + 4$ ,  $y(0) = 1$ ,  $y'(0) = 1$
- b) Find the orthogonal trajectories for the family of curves  $r^n \sin n\theta = a^n$ . [5+5]
- OR
- 3.a) In an L-R circuit an e.m.f. of  $10 \sin t$  volts is applied. If  $I(0) = 0$ , find the current  $I(t)$  in the circuit at any time  $t$ .
- b) Solve the Following differential equation  $y'' + 2y' + 5y = 4e^{-t} \cos 2t$ ,  $y(0) = 1$ ,  $y'(0) = 0$ . [5+5]

4.a) Find an LU factorization for the matrix  $\begin{bmatrix} 1 & 2 \\ -3 & -1 \end{bmatrix}$

b) In the following equations determine, for what value of "k" if any will the systems have  
i) unique solution ii) no solution iii) Infinitely many solutions

$$kx + 2y = 3$$

$$2x - 4y = -6$$

[5+5]

OR

5.a) Use either the Gaussian Elimination or the Gauss Jordan method to solve

$$x + 2y - 3z = 9$$

$$2x - y + z = 0$$

$$4x - y + z = 4$$

b) Using the theory of matrices, find the point such that the line of intersection of the planes

$$3x + 2y + z = -1 \text{ and } 2x - y + 4z = 5 \text{ cuts the plane } x + y + z = 4.$$

[5+5]

6.a) Obtain the Eigen values of the following matrix  $A = \begin{bmatrix} 2 & 0 & 1 \\ 0 & 2 & 0 \\ 1 & 0 & 2 \end{bmatrix}$  and verify whether its Eigen vectors are orthogonal.

b) Show that 0 is an Eigen value of a matrix A if and only if it is singular. [5+5]

OR

7.a) If  $A = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$  then show that  $A^n = A^{n-2} + A^2 - I$  for  $n \geq 3$ . Hence find  $A^{50}$ .

b) Show that the matrix  $A = \begin{bmatrix} 2 & 3 & 4 \\ 0 & 2 & -1 \\ 0 & 0 & 1 \end{bmatrix}$  is not similar to a diagonal matrix. [5+5]

8.a) If  $\sin u = \frac{x^2 y^2}{x+y}$  show that  $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = 3 \tan u$ .

b) If  $f(0) = 0$  and  $f'(x) = \frac{1}{1+x^2}$  then using Jacobians show that  $f(x) + f(y) = f\left(\frac{x+y}{1-xy}\right)$ . [5+5]

OR

9.a) Expand  $e^x \cos y$  in powers of  $x$  and  $\left(y - \frac{\pi}{2}\right)$ .

b) Show that the rectangular solid of maximum volume that can be inscribed in a given sphere is a cube. [5+5]

10. Find the general integrals of the linear partial differential equations

a)  $y^2 p - xy q = x(z - 2y)$

b)  $(y + zx)p - (x + yz)q = x^2 - y^2$ .

[5+5]

OR

11. Find complete integrals of the following equations

a)  $p+q=pq$

b)  $p^2 q(x^2 + y^2) = p^2 + q$ .

[5+5]

---ooOoo---